

# Should foods and beverages with lower nutrient density (i.e., those with added sugars, saturated fat, and sodium) contribute to item clusters, representative foods, and therefore the nutrient profiles for each food group and subgroup used in modeling the USDA Dietary Patterns?: Food Pattern Modeling Protocol

Christopher A. Taylor, PhD, RDN, LD, FAND,<sup>a,b</sup> Steven A. Abrams, MD,<sup>a,c</sup> Sarah Booth, PhD,<sup>a,d</sup> Carol Byrd-Bredbenner, PhD, RD, FAND,<sup>a,e</sup> Heather A. Eicher-Miller, PhD,<sup>a,f</sup> Teresa Fung, ScD, RD,<sup>a,g</sup> Valarie Blue Bird Jernigan, DrPH, MPH,<sup>a,h</sup> Sameera Talegawkar, PhD,<sup>a,i</sup> Deirdre Tobias, ScD,<sup>a,j</sup> Meghan Adler, MS, RDN,<sup>k</sup> Colleen Cruz, MPH, RDN,<sup>k</sup> Janet de Jesus, MS, RD,<sup>l</sup> Dana DeSilva, PhD, RD,<sup>m</sup> Laural Kelly English, PhD,<sup>n</sup> Stephenie Fu,<sup>o</sup> Hazel Hiza, PhD,<sup>k</sup> Kevin Kuczynski, MS, RD,<sup>k</sup> Emily Madan, PhD,<sup>p</sup> Verena McClain, MSc,<sup>p</sup> TusaRebecca Pannucci, PhD, MPH, RD,<sup>q</sup> Ramkripa Raghavan, DrPH, MPH, MSc,<sup>n</sup> Joe Rorabaugh-Irwin, MS, RD,<sup>m</sup> Kelley Scanlon, PhD, RD,<sup>r</sup> Eve Stody, PhD<sup>s</sup>

<sup>a</sup> Food Pattern Modeling and Data Analysis Subcommittee, 2025 Dietary Guidelines Advisory Committee

<sup>b</sup> The Ohio State University, Subcommittee Chair, Food Pattern Modeling

<sup>c</sup> University of Texas at Austin

<sup>d</sup> Tufts University, Committee Chair

<sup>e</sup> Rutgers, The State University of New Jersey

<sup>f</sup> Purdue University, Subcommittee Chair, Data Analysis

<sup>g</sup> Simmons University

<sup>h</sup> Oklahoma State University

<sup>i</sup> The George Washington University

<sup>j</sup> Harvard University

<sup>k</sup> Food Pattern Modeling Analyst; Nutrition Guidance and Analysis Division (NGAD), Center for Nutrition Policy and Promotion (CNPP), Food and Nutrition Service (FNS), U.S. Department of Agriculture (USDA)

<sup>l</sup> Designated Federal Officer and Nutrition Advisor, Office of Disease Prevention and Health Promotion (ODPHP); Office of the Assistant Secretary for Health (OASH), U.S. Department of Health and Human Services (HHS)

<sup>m</sup> Food Pattern Modeling Analyst, ODPHP; OASH, HHS

<sup>n</sup> Systematic Review Analyst, Nutrition Evidence Systematic Review (NESR) Branch; NGAD, CNPP, FNS, USDA

<sup>o</sup> Senior Policy Advisor; Deputy Administrator's Office; CNPP, FNS, USDA

<sup>p</sup> Food Pattern Modeling Analyst, Panum Telecom, under contract with FNS, USDA

<sup>q</sup> Branch Chief, Nutrition and Economic Analysis Branch (NEAB); NGAD, CNPP, FNS, USDA

<sup>r</sup> Senior Analytical Advisor; CNPP, FNS, USDA

<sup>s</sup> Director, NGAD; CNPP, FNS, USDA



**Suggested citation:** Taylor CA, Abrams SA, Booth S, Byrd-Bredbenner C, Eicher-Miller HA, Fung T, Jernigan VB, Talegawkar S, Tobias D, Adler M, Cruz C, de Jesus J, DeSilva D, English LK, Fu S, Hiza H, Kuczynski K, Madan E, McClain V, Pannucci T, Raghavan R, Rorabaugh-Irwin J, Scanlon K, Stoody E. *Should foods and beverages with lower nutrient density (i.e., those with added sugars, saturated fat, and sodium) contribute to item clusters, representative foods, and therefore the nutrient profiles for each food group and subgroup used in modeling the USDA Dietary Patterns? Food Pattern Modeling Protocol.* May 2023. U.S. Department of Agriculture, Food and Nutrition Service, Center for Nutrition Policy and Promotion, Nutrition Guidance and Analysis Division. Available at: <https://www.dietaryguidelines.gov/>

The contents of this document may be used and reprinted without permission. Endorsements by NGAD, CNPP, FNS, or USDA of derivative products developed from this work may not be stated or implied.

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons using assistive technology should be able to access information in this report. For further assistance please email [SM.FN.DIETARYGUIDELINES@USDA.gov](mailto:SM.FN.DIETARYGUIDELINES@USDA.gov).

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotope, American Sign Language, etc.) should contact the responsible agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at [How to File a Program Discrimination Complaint](#) and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by:

- (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410;
- (2) (2) fax: (202) 690-7442; or
- (3) (3) email: [program.intake@usda.gov](mailto:program.intake@usda.gov).

USDA is an equal opportunity provider, employer, and lender.

# Table of contents

---

<b>Table of contents</b> .....	<b>3</b>
<b>Rationale</b> .....	<b>4</b>
<b>Introduction</b> .....	<b>4</b>
<b>Methods</b> .....	<b>7</b>
Develop a protocol .....	7
Develop an analytic framework .....	8
Develop an analytic plan .....	11
Conduct analyses.....	13
Synthesize analyses.....	13
Develop conclusion .....	14
Recommend future research.....	14
Protocol amendments .....	14
<b>References</b> .....	<b>14</b>
<b>Acknowledgments and funding</b> .....	<b>15</b>
 Figure 1. Calculation of a weighted average nutrient profile for the Red and Orange Vegetables subgroup .....	 6
 Table 1. Age-sex groups included in analyses .....	 11
Table 2. Nutritional goals for analyses .....	12
Table 3. Protocol amendments.....	14

## Rationale

---

Food Pattern Modeling (FPM) is a methodology used to a) illustrate how hypothetical changes to the amounts or types of foods and beverages in a dietary pattern might affect meeting nutrient needs, and b) assist in defining quantitative dietary patterns that reflect the evidence for health-promoting diets synthesized from systematic reviews, while meeting energy and nutrient needs. Each food group and subgroup within the USDA Dietary Patterns has a defined nutrient profile that serves as the foundation of any FPM analysis. The nutrient profiles are calculated as the average of nutrient-dense forms of foods and beverages assigned to each food group or subgroup, weighted according to the proportions of reported consumption (by gram weight consumed) in the U.S. population.<sup>1</sup> Historically, nutrient profile calculations included the proportional contribution of intake relative to *all* foods and beverages in a food group or subgroup reported in *What We Eat in America*, National Health and Nutrition Examination Survey (WWEIA, NHANES). Although nutrient profiles of food groups and subgroups included foods and beverages regardless of their nutrient density in the composite to calculate the proportions of reported consumption (in grams), only nutrient-dense representative foods are used to estimate energy and nutrients for each food group or subgroup in calculating the nutrient profiles. This approach generates nutrient profiles that preferentially reflect foods and beverages with little added sugars, saturated fat, and sodium.

The following FPM analyses will derive new nutrient profiles for each food group and subgroup that remove foods and beverages lower in nutrient density (i.e., those with added sugars, saturated fat, and sodium) from these calculations. Of note, the revised approach is not suggesting that foods and beverages lower in nutrient density no longer contribute to a given food group or subgroup as defined in the USDA Food Patterns Equivalents Database.<sup>2</sup> Additional analyses will examine scenarios where foods and beverages with lower nutrient density may be added to an overall health promoting dietary pattern that includes recommended limits for energy, added sugars, saturated fats, and sodium.

## Introduction

---

To prepare for the development of the *Dietary Guidelines for Americans, 2025-2030*, the U.S. Departments of Health and Human Services (HHS) and Agriculture (USDA) identified a proposed list of scientific questions based on relevance, importance, potential impact to federal programs, and avoiding duplication, which were posted for public comment.<sup>1,3</sup> The Departments appointed the 2025 Dietary Guidelines Advisory Committee (Committee) in January 2023 to review evidence on the scientific questions. Their review forms the basis of their independent, science-based advice and recommendations to HHS and USDA, which is considered as the Departments develop the next edition of the *Dietary Guidelines*. These questions were refined and prioritized by the Committee for consideration in their review of the evidence.

The Committee will be asked to answer the following question using FPM analyses:

Considering each life stage, should changes be made to the USDA Dietary Patterns (Healthy U.S.-Style, Healthy Mediterranean-Style, and/or Healthy Vegetarian); should additional Dietary Patterns be developed/proposed based on:

- Findings from systematic reviews, data analysis, and/or FPM analyses; and
- Population norms (e.g., starchy vegetables are often consumed interchangeably with grains), preferences (e.g., emphasis on one staple grain versus another), or needs (e.g., lactose intolerance) of the diverse communities and cultural foodways within the U.S. population?

Changes to USDA Dietary Patterns may include modification to the amounts of food groups/subgroups and/or recategorization of food groups/subgroups, as well as subsequent changes to energy available for other uses, including for added sugars.

As part of that process and to address the overarching FPM question, the following question for analysis has been identified:

**Should foods and beverages with lower nutrient density (i.e., those with added sugars, saturated fat, and sodium) contribute to item clusters, representative foods, and therefore the nutrient profiles for each food group and subgroup used in modeling the USDA Dietary Patterns?**

The Committee will use FPM analyses to address this question, with support from USDA's FPM methods team. This protocol will establish methods used to develop and compare revised nutrient profiles to nutrient profiles developed using existing methods for each food group (i.e., Vegetables; Fruits; Grains; Dairy and Fortified Soy Alternatives; and Protein Foods) and subgroup (i.e., Dark-Green Vegetables; Red and Orange Vegetables; Beans, Peas, and Lentils; Starchy Vegetables; Other Vegetables; Whole Grains; Refined Grains; Meats, Poultry, and Eggs; Seafood; Nuts, Seeds, and Soy Products). Vegetables; Fruits; Grains; Dairy and Fortified Soy Alternatives; and Protein Foods) and subgroup (i.e., Dark-Green Vegetables; Red and Orange Vegetables; Beans, Peas, and Lentils; Starchy Vegetables; Other Vegetables; Whole Grains; Refined Grains; Meats, Poultry, and Eggs; Seafood; Nuts, Seeds, and Soy Products).

Historically, all foods and beverages reported in WWEIA, NHANES that contribute to a food group and/or subgroup are used in the development of nutrient profiles. The existing steps used to calculate a weighted average nutrient profile are described in detail below and visualized in Figure 1. An approach to revise these steps follows the description of existing approaches.

1. First, all foods and beverages, including combination foods and mixed dishes, reported in WWEIA, NHANES are disaggregated into ingredients and categorized into ~400 item clusters.<sup>4</sup> An item cluster is a group of similar foods or beverages (e.g., all sources of cooked carrots grouped together). Each food group and subgroup nutrient profile is comprised of a collection of item clusters, each representing the various types and sources of foods and beverages comprising a food group or subgroup, such as the Red and Orange Vegetables subgroup.
2. Second, each item cluster is assigned a single, nutrient-dense representative food to exemplify the energy and nutrient values that represent the item cluster in FPM. The *Dietary Guidelines for Americans, 2020-2025* defines nutrient-dense foods and beverages as those which provide vitamins, minerals, and other health-promoting components and have no or limited added sugars, saturated fat, and sodium. For FPM, a nutrient-dense representative food is chosen by selecting a single food or beverage within the item cluster with the least amounts of added sugars, saturated fats, and/or sodium (e.g., cooked carrots with no added sugars, fat, or salt to represent all sources of cooked carrots). In some cases, if there is no nutrient-dense food within an item cluster (e.g., flavored milk item clusters), then a nutrient-dense form of a food from a similar item cluster (e.g., unflavored skim milk) is selected as the representative food.
3. Third, dietary recall data from WWEIA, NHANES is used to calculate the proportion of consumption that each item cluster contributes to the entire food group or subgroup. The result is a composite for each food group or subgroup (i.e., the proportional contribution of each item cluster within each food group or subgroup). While the energy and nutrients assigned to each item cluster are represented by a nutrient dense food, the proportion of consumption of that item cluster to the entire food group or subgroup considers all food or beverage sources, regardless of nutrient density (e.g., consumption of all cooked carrots regardless of if they contain added sugars, salt, and/or fat or are used as an ingredient in a food (i.e., carrots in carrot cake) to calculate the proportional contribution of the item cluster, "Carrots, cooked" to the subgroup "Red and Orange Vegetables").


- Fourth, a weighted average nutrient profile is calculated to develop the nutrient profile for each food group and subgroup. The calculation of each nutrient profile considers the proportional contribution from multiple item clusters. The percent contribution of each item cluster within a food group or subgroup is multiplied by the energy and nutrients in their assigned nutrient-dense representative foods. The results are summed to determine the overall nutrient profile for each food group and subgroup. The following formula is used to calculate each nutrient profile:

$$\sum (\% \text{ contribution of each item cluster} \times \text{nutrients in the representative food})$$

The 2020 Dietary Guidelines Advisory Committee used FPM to explore the development of dietary patterns for individuals less than 2 years of age.<sup>5</sup> The nutrient profiles for this age group were developed using the same steps, but with a few distinctions. The foods and beverages were limited to those reported (by proxy reporting) for this age group (i.e., < 2 years of age) and the proportion of consumption of each item cluster was specific to proportions of consumption in this age group. Differences in the representative foods used, compared to those used for nutrient profiles in FPM for ages 2 and older, were the following:

- Whole milk was used instead of fat-free milk
- Reduced-fat plain yogurt was used instead of fat-free yogurts (plain or flavored with non-caloric sweeteners)
- Reduced-fat cheeses were used as representative foods for all cheese item clusters instead of using skim or fat-free cheese options when available

**Figure 1. Calculation of a weighted average nutrient profile for the Red and Orange Vegetables subgroup**



\*2020 FPM Report 2+: [https://www.dietaryguidelines.gov/sites/default/files/2020-07/FoodPatternModeling\\_Report\\_2YearsandOlder.pdf](https://www.dietaryguidelines.gov/sites/default/files/2020-07/FoodPatternModeling_Report_2YearsandOlder.pdf)

Each nutrient profile includes energy, the macronutrients, 12 vitamins, and 8 minerals. For example, the nutrient profile for the Red and Orange Vegetables subgroup includes ~45 kcal, ~2.4 g of fiber, and ~43 mg

of vitamin C, which reflects the weighted average contribution of the nutrients in the nutrient-dense representative foods that represent each item cluster that make up Red and Orange Vegetables subgroup.

This protocol describes a multi-phased approach to exclude foods and beverages that are lower in nutrient density. The same general steps for constructing nutrient profiles remain; however, instead of including all foods reported in WWEIA, NHANES, foods and beverages identified as being lower in nutrient density will be excluded from the calculation of the revised nutrient profile. Of note, the revised approach is not suggesting that foods and beverages lower in nutrient density no longer contribute to a given food group or subgroup as defined in the USDA Food Patterns Equivalents Database. Rather, some foods and beverages may be excluded from the calculation of a nutrient-dense nutrient profile for each food group or subgroup.

The development of a nutrient profile allows the Committee to estimate the anticipated nutrient composition for each food group and subgroup that could be obtained by eating a variety of foods from each group in nutrient-dense forms, while further limiting the inclusion of foods and beverages that have saturated fat, added sugars, and sodium. The resulting nutrient profiles can then be used in analyses to assess nutrient adequacy of dietary patterns for various individuals. The Committee will consider if the existing approach or a revised approach to calculating nutrient profiles will be used for subsequent FPM analyses.

## Methods

---

This section presents an overview of the methods, or the process, that will be used by the Committee to answer the question:

**Should foods and beverages with lower nutrient density (i.e., those with added sugars, saturated fat, and sodium) contribute to item clusters, representative foods, and therefore the nutrient profiles for each food group and subgroup used in modeling the USDA Dietary Patterns?**

## Develop a protocol

A FPM protocol is the plan for how USDA's FPM methodology will be used to conduct specific FPM analyses. The protocol is established by the Committee before the analysis is conducted. The protocol describes the components of the FPM process, including the analytic framework, analytic plan, analysis synthesis, conclusion development, and future research recommendations. It is developed through Committee discussion of the strengths and limitations for various analysis types and exercises to identify the most appropriate and relevant methods to answer each FPM question. FPM is an iterative process, thus results from initial analyses may inform refinement of this protocol or subsequent protocols for other research questions.

When reviewing questions or topics addressed by prior Committees, the Committee uses the previous analytic framework, plan, and protocol to inform and refine their current approaches. Any changes to this protocol will be described in **Table 3. Protocol amendments**.

## Develop an analytic framework

An analytic framework represents the overall scope of the FPM analyses, including the population, type of analyses, and data sources identified to answer the question. It also includes the definitions of key terms.

---

### Question:

**Should foods and beverages with lower nutrient density (i.e., those with added sugars, saturated fat, and sodium) contribute to item clusters, representative foods, and therefore the nutrient profiles for each food group and subgroup used in modeling the USDA Dietary Patterns?**

---

### Population:

The nutrient profiles tested in these FPM analyses are based on dietary intake data among the U.S. population ages 1 year and older.

The nutrient profiles tested in these FPM analyses will be applied to the existing [U.S. Healthy-Style Dietary Patterns](#) published for ages 12 through 23 months and for ages 2 years and older.<sup>1</sup>

### Types of Analyses:

The overall FPM methodology used to develop and update the USDA Food Patterns includes: (1) identifying appropriate energy levels for the patterns; (2) identifying nutritional goals for the patterns; (3) establishing food groupings and food group amounts; (4) determining the amounts of energy and nutrients that would be provided by consuming various foods within each food group or subgroup; and (5) evaluating nutrient levels in each pattern against nutritional goals. Finally, (6) multiple iteration and re-evaluation of revised nutrient profiles may be required to test differences in the exclusions of foods and beverages that are lower in nutrient density from being used to calculate nutrient profiles.

**This question will focus on the step 4, determining the amounts of energy and nutrients that would be obtained by consuming various foods within each food group.** In this step, nutrient profiles are calculated for each food group and subgroup. Analyses for the protocol will test the implications of using revised nutrient profiles that exclude foods and beverages based on nutrient density (i.e., added sugars, saturated fat, and sodium content) compared to existing nutrient profiles that include all food and beverage sources of a food group or subgroup. Revised nutrient profiles can then be applied in step 5 to evaluate implications for nutrient levels in each pattern.

FPM analyses for answering these questions will involve:

- Calculate a nutrient profile for each food group and subgroup using existing methods that does not exclude the contribution of identified less nutrient-dense foods and beverages.
- Identify foods and beverages that are lower in nutrient density and that currently contribute to item clusters, representative foods, and, therefore, nutrient profiles for each food group and subgroup used in modeling the Healthy U.S.-Style Dietary Pattern.
- Calculate a revised nutrient profile for each food group and subgroup that excludes the contribution of identified foods and beverages that are lower in nutrient density.
- Compare food group and subgroup nutrient profiles that exclude the contribution of foods and beverages that are lower in nutrient density to nutrient profiles calculated using existing methods that do not exclude less nutrient-dense foods and beverages.
  - Evaluate proportional contribution of item clusters (i.e., each item cluster's percent contribution) to the total composite of each food group and subgroup.



- Compare total energy and nutrients provided in the Healthy U.S.-Style Dietary Pattern, as defined in the *Dietary Guidelines for Americans, 2020-2025* for ages 12 through 23 months and for ages 2 years and older, when nutrient profiles are calculated using the revised methods to the total energy and nutrients provided when nutrient profiles are calculated using existing methods.<sup>1</sup>
  - Identify the age-sex groups and life stages for whom nutrient needs are met or not met at each calorie level.
- FPM is an iterative process (i.e., repeated analyses), thus results from initial analyses may inform refinement of this protocol. Any changes to this protocol will be described in **Table 3. Protocol amendments**.
- Determine if existing or revised nutrient profiles will be used for subsequent FPM analyses.

## Data Sources:

- What We Eat in America, NHANES 2017-2018, individuals 1 years and over, days 1 and 2, weighted to produce nationally representative estimates. Available: [ars.usda.gov/nea/bhnrc/fsrg](https://ars.usda.gov/nea/bhnrc/fsrg)
- FNDDS 2017-2018: U.S. Department of Agriculture, Agricultural Research Service. 2020. *USDA Food and Nutrient Database for Dietary Studies 2017-2018*. Available: [ars.usda.gov/nea/bhnrc/fsrg](https://ars.usda.gov/nea/bhnrc/fsrg)
- U.S. Department of Agriculture, Agricultural Research Service, Beltsville Human Nutrition Research Center, Food Surveys Research Group, Beltsville, Maryland, Food Patterns Equivalents Databases and Datasets 2017-2018. Available: [ars.usda.gov/nea/bhnrc/fsrg](https://ars.usda.gov/nea/bhnrc/fsrg)
- FPED 2017-2018: Bowman SA, Clemens JC, Friday JE, and Moshfegh AJ. 2020. Food Patterns Equivalents Database 2017-2018: Methodology and User Guide [Online]. Food Surveys Research Group, Beltsville Human Nutrition Research Center, Agricultural Research Service, U.S. Department of Agriculture, Beltsville, Maryland. October 2020. Available: [ars.usda.gov/nea/bhnrc/fsrg](https://ars.usda.gov/nea/bhnrc/fsrg)
- U.S. Department of Agriculture, Agricultural Research Service. 2020. What We Eat in America Food Categories 2017-2018. Available: [ars.usda.gov/nea/bhnrc/fsrg](https://ars.usda.gov/nea/bhnrc/fsrg)
- SR 28: US Department of Agriculture, Agricultural Research Service. 2016. Nutrient Data Laboratory. USDA National Nutrient Database for Standard Reference, Release 28 (Slightly revised). Version Current: May 2016. Available: [ars.usda.gov/nea/bhnrc/mafcl](https://ars.usda.gov/nea/bhnrc/mafcl)
- U.S. Department of Agriculture and U.S. Department of Health and Human Services. *Dietary Guidelines for Americans, 2020-2025*. 9th Edition. December 2020. Available at [DietaryGuidelines.gov](https://www.dietaryguidelines.gov)
- National Academies of Sciences, Engineering, and Medicine. 2023. *Dietary Reference Intakes for Energy*. Washington, DC: The National Academies Press. <https://doi.org/10.17226/26818>
- National Academies of Sciences, Engineering, and Medicine. 2019. *Dietary Reference Intakes for Sodium and Potassium*. Washington, DC: The National Academies Press. doi: <https://doi.org/10.17226/25353>
- National Academies of Sciences, Engineering, and Medicine. 2011. *Dietary Reference Intakes for Calcium and Vitamin D*. Washington, DC: The National Academies Press. <https://doi.org/10.17226/13050>
- Institute of Medicine. 2006. *Dietary Reference Intakes: The Essential Guide to Nutrient Requirements*. Washington, DC: The National Academies Press. <https://doi.org/10.17226/11537>
- Fryar CD, Carroll MD, Gu Q, Afful J, Ogden CL. Anthropometric reference data for children and adults: United States, 2015–2018. National Center for Health Statistics. Vital Health Stat 3(46). 2021. Available: [cdc.gov/nchs/nhanes](https://cdc.gov/nchs/nhanes)

## Key definitions:

Note: Key definitions in this protocol include existing definitions used by the 2020 Dietary Guidelines Advisory Committee and/or published in the *Dietary Guidelines for Americans, 2020-2025*, such as definitions operationalized in the Healthy U.S.-Style Dietary Pattern.<sup>1, 6</sup> The 2025 Dietary Guidelines Advisory Committee will continue to consider terminology and implications of terms related to health equity and/or communication to the public. Future revisions to existing definitions will be noted.

**Food Groups and Subgroups:** The Healthy U.S.-Style Dietary Pattern provides amounts of five major food groups and subgroups including:

- Fruits
- Vegetables:
  - *Dark-Green; Red and Orange; Beans, Peas, and Lentils; Starchy; and Other*
- Dairy and Fortified Soy Alternatives
- Grains:
  - *Whole Grains and Refined Grains*
- Protein Foods:
  - *Meats, Poultry, and Eggs; Seafood; Nuts, Seeds, and Soy Products*

\*For this protocol, Beans, Peas, and Lentils will only be modeled in the vegetable group.

**Oils:** Oils are sources of essential fatty acids and include canola, corn, olive, peanut, safflower, soybean, and sunflower oils. Oils also are naturally present in nuts, seeds, seafood, olives, and avocados. The fat in some tropical plants, such as coconut oil, palm kernel oil, and palm oil, are not included in the oils category because they contain a higher percentage of saturated fat than do other oils.

**Added sugars:** Added sugars are either added during the processing of foods or are packaged as sweeteners (e.g., a bag of table sugar). Added sugars include sugars (free, mono- and disaccharides), sugars from syrups and honey, and sugars from concentrated fruit or vegetable juices that are in excess of what would be expected from the same volume of 100 percent fruit or vegetable juice of the same type.<sup>7</sup> Naturally occurring sugars, such as those in fruit or milk, are not defined as added sugars. Specific examples of added sugars that can be listed as an ingredient include brown sugar, corn sweetener, corn syrup, dextrose, fructose, glucose, high-fructose corn syrup, honey, invert sugar, lactose, malt syrup, maltose, molasses, raw sugar, sucrose, trehalose, and turbinado sugar.

**Limits on calories for other uses (as defined in the Healthy U.S.-Style Dietary Pattern)<sup>1, 6</sup>:** Foods are assumed to be in nutrient-dense forms, which are lean or low-fat and prepared with minimal added saturated fat, added sugars, refined starches, or sodium. If all food choices to meet food group recommendations are in nutrient-dense forms, a small number of calories remain within the overall limit of the pattern (i.e., limit on calories for other uses). The amount of calories depends on the total calorie level of the pattern and the amounts of food from each food group required to meet nutritional goals. Calories up to the specified limit can be used for added sugars, refined starches, saturated fat, and/or alcohol (for nonpregnant adults of legal drinking age only), or to eat more than the recommended amount of food in a food group.

**Item Clusters:** Identified groupings of the same or similar foods or beverages that make up each food group and subgroup.

**Nutrient-Dense Representative Foods:** For the purpose of USDA's FPM, each item cluster is assigned a nutrient-dense representative food which are those foods or beverages that represent the forms with the least amounts of added sugars, sodium, and saturated fats. The nutrient composition of the nutrient-dense representative food is used to represent the nutrient composition of the entire item cluster when calculating the nutrient profile for a food group or subgroup.

**Nutrient Profiles:** The proportional nutrient composition from the item clusters that represent each food group and subgroup from the variety of foods in each food group in their nutrient-dense forms. The nutrient profiles are based on a weighted average of nutrient-dense forms of foods (i.e., a composite of nutrient-dense forms of foods and beverages within a food group or subgroup). The weighted average calculation considers a range of food choices in the United States, but in nutrient-dense forms and results in a food pattern that can be adapted to fit an individual's preferences.

## Develop an analytic plan

### Establish energy levels:

Dietary Reference Intakes (DRI) formulas are used to calculate Estimated Energy Requirements (EER) for each age-sex group, with three age groups specific to pregnancy and lactation: 14-18 years, 19-30 years and 31-50 years.<sup>8</sup> (See Table 1.) EER is based on sex, age, height, weight, and physical activity level. Median body height and weight for normal body mass index (BMI) are used to calculate appropriate energy levels for each age-sex group.<sup>1</sup> The EERs for pregnancy account for additional energy expenditure and deposition that includes the products of conception and accretion of maternal tissues; those for lactation account for energy expenditure associated with human milk output and weight loss. The dietary patterns are not specific to individual age-sex groups or life stages, with the exception of patterns intended specifically for individuals 12 through 23 months ranging from 700 to 1,000 kcal in 100 kcal intervals, and a second set of patterns intended for ages 2 years and older that includes 12 energy levels from 1,000 to 3,200 kcal at 200 kcal “step” intervals intended to cover energy needs for the majority of the population.

**Table 1. Age-sex groups included in analyses**

Children (Male/Female)	Males	Females	Pregnancy (Per 1 <sup>st</sup> , 2 <sup>nd</sup> , and 3 <sup>rd</sup> Trimesters)	Lactation (Per 0-6 and 7-12 months postpartum)
1-3 years	4-8 years	4-8 years	14-18 years	14-18 years
	9-13 years	9-13 years	19-30 years	19-30 years
	14-18 years	14-18 years	31-50 years	31-50 years
	19-30 years	19-30 years		
	31-50 years	31-50 years		
	51+ years	51+ years		

### Establish nutritional goals:

Specific nutritional goals for each dietary intake pattern (i.e., by energy level) are selected based on the DRI age-sex group(s) for which the pattern is appropriate. If a dietary intake pattern at an energy level aims to meet the needs for more than one age-sex group, the pattern is evaluated against the nutrient goals for all those groups. Goals for energy, 3 macronutrients, 3 fatty acids, 12 vitamins, 8 minerals, added sugars, and fiber are based on DRI reports released between 1997 and 2023 and on quantitative recommendations in the current *Dietary Guidelines for Americans, 2020-2025* (DGA).<sup>1, 8-11</sup> The macronutrients, fatty acids, vitamins, and minerals are specified in Table 2. Because the dietary patterns are designed as a framework for an individual to achieve a healthy dietary pattern, the goals are the Recommended Dietary Allowance (RDA) amounts for nutrients having an RDA. The Adequate Intake (AI) is used when an RDA is not published. The lowest energy level (for sedentary individuals, determined in step 1) rounded to the nearest 200 kcal energy level and associated pattern is determined for each age-sex group and used in evaluating the patterns against nutritional goals.

**Table 2. Nutritional goals for analyses**

<b>Food Component</b>	<b>Specific Nutrients (and Source of Goal <sup>a</sup>)</b>
Energy	Energy (EER)
Macronutrients	Carbohydrate (AMDR/RDA), Protein (AMDR/RDA), Total Lipid (AMDR)
Fatty acids	Saturated Fatty Acids (DGA), 18:2 Linoleic Acid (AI), 18:3 Linolenic Acid (AI)
Vitamins	Vitamin A (RDA), Vitamin C (RDA), Vitamin D (RDA), Vitamin E (RDA), Vitamin K (AI), Thiamin (RDA), Riboflavin (RDA), Niacin (RDA), Vitamin B6 (RDA), Folate (RDA), Vitamin B12 (RDA), Choline (AI)
Minerals	Calcium (RDA), Copper (RDA), Iron (RDA), Magnesium (RDA), Phosphorus (RDA), Potassium (AI), Sodium (CDRR), Zinc (RDA)
Added Sugars	Added Sugars (DGA, <10% of kcal)
Fiber	Total Dietary Fiber (14g/1,000 kcal)

<sup>a</sup> AI = Adequate Intake, AMDR = Acceptable Macronutrient Distribution Range, CDRR = Chronic Disease Risk Reduction Level, DGA = *Dietary Guidelines for Americans, 2020-2025*, RDA = Recommended Dietary Allowance

**Establish food groupings and amounts:**

Existing food groups and subgroups in the USDA Healthy U.S.-Style Dietary Pattern published in the *Dietary Guidelines for Americans, 2020-2025* are used in these analyses.<sup>1</sup> The food group and subgroup amounts are informed by the Healthy U.S.-Style Dietary Pattern for ages 12 through 23 months and for ages 2 years and older.

**Determine the amounts of nutrients that would be obtained by consuming various foods within each group:**

- A composite system is used to determine the anticipated energy and nutrient content, or nutrient profile, of each food group or subgroup as described below.
  - All foods reported by individuals ages 1 year and older as part of WWEIA, NHANES 2017-2018 are disaggregated into their ingredients.<sup>12</sup>
    - Existing nutrient profile development approach: all foods and beverages that have a food group or subgroup contribution are included in the set of foods used to calculate nutrient profiles.
    - Revised nutrient profile development approach: some foods and beverages that are lower in nutrient density are excluded from the set of foods used to calculate nutrient profiles.
      - Exclude foods/beverages based on WWEIA Food Categories and companion item clusters.
      - Exclude foods/beverages for which less than [defined proportion] of the total ingredients contribute to a food group/subgroup.
      - Exclude item clusters when a nutrient-dense representative food would not be a practical, nutrient-dense alternative for the foods and beverages within an item cluster.

- Exclude item clusters when the representative food is an outlier compared to other nutrient-dense representative foods based on the amount of added sugars, saturated fat, and/or sodium.
  - Similar ingredients are aggregated into food item clusters.
  - A nutrient-dense form of the food is selected as the representative food for each cluster.
  - The proportional intake of each item cluster within each food group or subgroup is calculated and used to compute a weighted average of nutrient-dense forms of foods representing each food item cluster.
    - The proportional intake is calculated for ages 1 year or older similar to methods from previous updates.<sup>1</sup>
- Using the nutrients in each representative food and the item cluster's proportional intake using the life stage approach (children less than 2 years and the population 2 years and older), a nutrient profile is calculated for each food group or subgroup. Thus, a nutrient profile specific to each child age 12 through 23 months and to the rest of the population is created and used to estimate the anticipated nutrients in the patterns. Nutrient profiles are also calculated for oils and solid fats using food supply data to determine proportional intakes.

## Evaluate nutrient level in each pattern against nutritional goals:

Using the revised nutrient profiles that apply to children less than 2 years and the population 2 years and older, the nutrients provided by amounts recommended in the *Dietary Guidelines for Americans, 2020-2025* from each food group (and oils) are compared to the age, sex, and life stage-specific goals (usually at least 90% of the RDA or AI).

## Iteration and re-evaluation:

After identifying the implications of the defined revised nutrient profiles and their comparison to the nutrient profiles calculated with existing methods, the Committee may use a stepwise, iterative approach to make adjustments. This may result in testing a different set of defined exclusions of foods and beverages and re-evaluation of the resulting nutrient profiles. Any changes to this protocol will be described in **Table 3. Protocol amendments**.

## Conduct analyses

The USDA FPM methods team, in collaboration with the Committee, will use the analytic framework and analytic plan as a guide for conducting analyses and preparing tables and reports describing the analytic results for each analysis.

The first level of analysis will be by population with results described for each age-sex groups and life stage (e.g., pregnancy and lactation). Depending on the available data, subsequent analyses may be based on population subgroups, race/ethnicity, and/or socioeconomic position.

## Synthesize analyses

The Committee will describe, compare, and combine the evidence from all FPM analyses conducted to answer this FPM question. Synthesis of the analyses will involve summarizing results with particular emphasis on implications for each life stage: infants, children, adolescents, adults, older adults, and individuals who are pregnant or lactating. Implications for each of the existing USDA Dietary Patterns or rationale for new pattern development will also be included.

The analyses related to each individual protocol will be considered together in answering the primary question.

## Develop conclusion

The Committee will review and discuss the synthesis of the analyses to develop conclusion statements for each FPM question. Conclusions from this protocol will be used along with conclusions from all other FPM protocols to collectively inform the Committee's advice on the development or refinement of healthy dietary patterns.

## Recommend future research

The Committee will identify and document research gaps and methodological limitations throughout the FPM process. These gaps and limitations will be used to develop research recommendations that describe the research, data, and methodological advances that are needed to strengthen the process to test and develop healthy dietary patterns. Rationales for the necessity of additional or stronger research may also be provided with the research recommendations.

## Protocol amendments

No amendments to the protocol have been made at this time. Any future amendments will be documented below in **Table 3**.

**Table 3. Protocol amendments**

Date	Protocol change	Description
N/A	N/A	N/A

## References

1. U.S. Department of Agriculture and U.S. Department of Health and Human Services. *Dietary Guidelines for Americans, 2020-2025, 9th Edition*. 2020. Accessed May 23, 2023.
2. U.S. Department of Agriculture, Agricultural Research Service. Food Patterns Equivalents Database. Updated May 4, 2023. Accessed May 23, 2023.
3. U.S. Department of Agriculture and U.S. Department of Health and Human Services. Learn About the Process. Accessed May 24, 2023.
4. 2020 Dietary Guidelines Advisory Committee and Food Pattern Modeling Team. *Added Sugars: Food Pattern Modeling: Ages 2 Years and Older*. Washington, DC: U.S. Department of Agriculture; 2020. *2020 Dietary Guidelines Advisory Committee Project*.
5. 2020 Dietary Guidelines Advisory Committee and Food Pattern Modeling Team. *Food Pattern Modeling: Under 2 Years of Age*. Washington, DC: U.S. Department of Agriculture; 2020. *2020 Dietary Guidelines Advisory Committee Project*.
6. Dietary Guidelines Advisory Committee. *Scientific Report of the 2020 Dietary Guidelines Advisory Committee: Advisory Report to the Secretary of Agriculture and the Secretary of Health and Human Services*. Washington, DC: U.S. Department of Agriculture, Agricultural Research Service; 2020.
7. U.S. Department of Health and Human Services, Food and Drug Administration. Industry Resources on the Changes to the Nutrition Facts Label: Added Sugars. Updated January 11, 2022. Accessed May 24, 2023.
8. National Academies of Sciences, Engineering, and Medicine. *Dietary Reference Intakes for Energy*. Washington, DC: The National Academies Press; 2023.
9. Institute of Medicine. *Dietary Reference Intakes: The Essential Guide to Nutrient Requirements*. Washington, DC: The National Academies Press; 2006.

10. National Academies of Sciences, Engineering, and Medicine. *Dietary Reference Intakes for Sodium and Potassium*. Washington, DC: The National Academies Press; 2019.
11. National Academies of Sciences, Engineering, and Medicine. *Dietary Reference Intakes for Calcium and Vitamin D*. Washington, DC: The National Academies Press; 2011.
12. U.S. Department of Agriculture, Agricultural Research Service. What We Eat in America Documentation and Data Sets. Updated August 8, 2022. Accessed May 24, 2023.

## Acknowledgments and funding

---

The Committee members are involved in: establishing all aspects of the protocol, which presents the plan for how they are planning to examine the scientific evidence, including the development of an analytic framework and analytic plan; synthesizing analysis results; and writing conclusion statements. The analytic framework and plan provide details about the types of analyses that will be conducted, synthesized, and from which conclusions will be drawn to inform subsequent FPM questions and the Committee's advice on the development or refinement of healthy dietary patterns. The FPM Methods Team, with assistance from Federal Liaisons and Project Leadership, supports the Committee by facilitating, executing, and documenting the work necessary.

**Funding:** United States Department of Agriculture, Food and Nutrition Service, Center for Nutrition Policy and Promotion, Alexandria, VA; Department of Health and Human Services, Office of Disease Prevention and Health Promotion, Rockville, MD.